

Common Fixed Point Theorems for Nonself – Mapping in Metrically Convex Spaces

Geeta Modi[†], Arvind Gupta[‡] and Varun singh^{§1}

[†]Head & Professor of Department Mathematics,
Govt. M.V.M. Bhopal, MP, India.
modi.geeta@gmail.com

[‡]Professor of Department Mathematics,
Govt. M.V.M. Bhopal, MP, India.
arvind533@yahoo.com

[§]Research Scholar,
Govt. M.V.M. Bhopal, MP, India.
varunsinghhh@gmail.com

Abstract : A common fixed point theorem for pair of nonself- mapping is proved in complete metrically convex metric space, which generalize earlier results due to M.Imdad [6] M.D. khan [4], MS Khan [5], Bianchini [12], Chatterjea [13], and others.

Keywords : Banach space, metrically convex space , fixed point.

1 Introduction

In many applications the mappings under examination may not always be self- mapping, therefore fixed point theorems for nonself-mapping from a natural subject for investigation. Assad and Kirk [8] initiated

¹Corresponding author E-Mail: varunsinghhh@gmail.com(Varun singh)
AMS Subject Classification: 30C15, 30C10, 26C10.

the study of fixed point for nonself –mappings in metrically convex spaces, noticed that with some kind of metric convexity, domain and range of mapping under examination can be considered of more varied type. In recent years this technique due to Assad and Kirk [8] has been utilized by many researchers of domain and by now there exists considerable literature on this topic. To mention a few, we cite [2, 3, 4, 5, 7, 8, 9, 10].

Assad [7] gave sufficient condition for nonself- mappings defined on a closed subset of complete metrically convex metric spaces satisfying Kannan type mappings [11] which have been generalized by M.S. Khan [5]. For the sake of completeness, we stat that main result of M.S. Khan [5].

Theorem 1.1. *Let (X, d) be a complete metrically convex space and K a closed nonempty subset of X . Let $T : K \rightarrow X$ be the mapping satisfying the inequality*

$$d(Tx, Ty) \leq C \max \{d(x, Tx), d(y, Ty)\} + C' \{d(x, Tx) + d(y, Ty)\} \quad (1.1)$$

for every x, y in K , where C and C' are non-negative reals such that

$$\begin{aligned} \max \left\{ \frac{C + C'}{1 - C'}, \frac{C'}{1 - C - C'} \right\} &= h > 0 \\ \max \left\{ \frac{1 + C + C'}{1 - C'} h, \frac{1 + C'}{1 - C - C'} h \right\} &= h' > 0 \\ \max \{h, h'\} &= h'' > 0 \end{aligned}$$

Further, for every x in ∂K , $Tx \in K$. Then T has a unique fixed point in K .

Currently M. Imdad and Ladlay Khan [6] generalized this result as,

Theorem 1.2. *Let (X, d) be a complete metrically convex metric space and K a closed nonempty subset of X . if F is generalized T contraction mapping of K into X satisfying the following:*

1. $\delta K \subseteq TK$, $FK \cap K \subseteq TK$
2. $Tx \in \delta K \Rightarrow Fx \in K$
3. (F, T) is coincidentally commuting
4. TK is closed in X .

Then F and T have a unique common fixed point.

Definition 1.3 ([6]). *Let (X, d) be a metric space and K a nonempty subset of X . Let $F, T : K \rightarrow X$ be a pair of maps which satisfy the condition*

$$\begin{aligned} \phi(d(Fx, Fy)) \leq a \max \left\{ \frac{1}{2} \phi(d(Tx, Ty)), \phi(d(Tx, Fx)), \phi(d(Ty, Fy)) \right\} \\ + b \{ \phi(d(Tx, Fy)) + \phi(d(Ty, Fx)) \} \end{aligned} \quad (1.2)$$

for all distinct $x, y \in K$, $a, b \geq 0$ such that $a + 4b < 1$ and let $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be an increasing continuous function for which the following properties hold:

$$\phi(t) = 0 \iff t = 0, \quad \phi(2t) \leq 2\phi(t)$$

Then F is called generalized T contraction mapping of K into X .

2 Preliminaries

We necessitate the following in the sequel

Definition 2.1 ([9]). Let K be a nonempty subset of a metric space (X, d) and $T : K \rightarrow X$. The pair (F, T) is said to be weakly commuting if for every $x, y \in K$ with $x = Fy$ and $Ty \in K$

$$(Tx, FTy) \leq d(Ty, Fy) \quad (2.1)$$

Definition 2.2 ([10]). Let K be a nonempty subset of a metric space (X, d) and $F, T : K \rightarrow X$. The pair (F, T) is said to be compatible if every sequence $\{x_n\} \subset K$ and from the relation

$$\lim_{n \rightarrow \infty} d(Fx_n, Tx_n) = 0. \quad (2.2)$$

And $Tx_n \in K$ (for every) $n \in \mathbb{N}$, it follows that

$$\lim_{n \rightarrow \infty} d(Ty_n, FTx_n) = 0 \quad (2.3)$$

for every sequence $y_n \in K$ such that $y_n = Fx_n$, $n \in \mathbb{N}$

Definition 2.3 ([3]). A pair of nonself-mapping (F, T) on a nonempty subset K of a metric space (X, d) is said to be coincidentally commuting if $Tx, Fx \in K$ and $Tx = Fx \implies FTx = TFx$

Definition 2.4 ([8]). A metric space (X, d) is said to be metrically convex if for any distinct $x, y \in X$, there exists a point $z \in X$ with $x \neq z \neq y$ such that

$$(x, z) + d(z, y) = d(x, y) \quad (2.4)$$

Lemma 2.5 ([8]). Let K be a nonempty closed subset of a metrically convex metric space X . if $x \in K$ and $z \notin K$, then there exists a point $z \in \delta K$ (the boundary of K) such that

$$d(x, z) + d(z, y) = d(x, y) \quad (2.5)$$

3 Our main result runs as follows

Theorem 3.1. Let (X, d) be a complete metrically convex metric space and K a closed nonempty subset of X . Let $F, T : K \rightarrow X$ be the mappings satisfying the inequality

$$(Fx, Fy) \leq (a + b) d(Fx, Tx) + b [\max \{d(Tx, Fx), d(Tx, Ty)\} + d(Ty, Fy)] \quad (3.1)$$

for every $x, y \in K$, where a, b, c are nonnegative reals such that

$$\frac{a+b+c}{1-c} = h > 0$$

and

$$(i) \delta K \subseteq TK, FK \cap K \subseteq TK$$

$$(ii) Tx \in \delta K \Rightarrow Fx \in K$$

(iii) (F, T) is coincidentally commuting

(iv) TK is closed in X

Then F and T have a unique common fixed point.

Proof. Firstly, we proceed to construct two sequences $\{x_n\}$ and $\{y_n\}$ in the following way: Let $x \in \delta K$. Then (due to $\delta K \subset TK$) there exists a point $x_0 \in K$ such that $x = Tx_0$. Since $Tx \in \delta K \implies Fx \in K$, one concludes that $Fx_0 \in FK \cap K \subseteq TK$. Let $x_1 \in K$ be such that $y_1 = Tx_1 = Fx_0$.

Let $y_2 = Fx_1$, if $y_2 \in K$, then $y_2 \in FK \cap K \subset TK$ which implies that there exists a point $x_2 \in K$ such that $y_2 = Tx_2$. If $y_2 \notin K$, then there exists a point $t \in \delta K$ such that

$$d(Tx, t) + d(t, y_2) = d(Tx_1, y_2) \quad (3.2)$$

Since $t \in \delta K \subseteq TK$ there exists a point $x_2 \in K$ such that $t = Tx_2$ so that

$$d(Tx_1, Tx_2) + d(Tx_2, y_2) = d(Tx_1, y_2) \quad (3.3)$$

Thus repeating the foregoing arguments one obtains two sequences $\{x_n\}$ and $\{y_n\}$ such that

$$(v) y_{n+1} = Fx_n$$

$$(vi) y_n \in K \Rightarrow y_n = Tx_n \quad \text{or} \quad y_n \notin K \Rightarrow Tx_n \in \delta K$$

and

$$d(Tx_{n-1}, Tx_n) + d(Tx_n, y_n) = d(Tx_{n-1}, y_n). \quad (3.4)$$

We denote

$$P = \{Tx_i \in \{Tx_n\} : Tx_i = y_i\}$$

$$Q = \{Tx_i \in \{Tx_n\} : Tx_i \neq y_i\}$$

Obviously, the two consecutive terms cannot lie in Q . Now, we distinguish the following three cases,

Case 1. If $Tx_n, Tx_{n+1} \in P$ then

$$\begin{aligned} d(Tx_n, Tx_{n+1}) &= d(y_n, y_{n+1}) \\ &= d(Fx_{n-1}, Fx_n) \\ &\leq (a+c)d(Fx_{n-1}, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, Fx_{n-1}), d(Tx_{n-1}, Tx_n)\} + d(Tx_n, Fx_n)] \\ &= (a+c)d(y_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, y_n), d(Tx_{n-1}, Tx_n)\} + d(Tx_n, y_{n+1})] \\ &= (a+c)d(Tx_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, Tx_n), d(Tx_{n-1}, Tx_n)\} + b d(Tx_n, Tx_{n+1})] \end{aligned}$$

$$\begin{aligned}
(1-b)d(Tx_n, Tx_{n+1}) &\leq (a+c)d(Tx_n Tx_{n-1}) + b d(Tx_{n-1}, Tx_n) \\
(1-b)d(Tx_n, Tx_{n+1}) &\leq (a+c+b)d(Tx_{n-1}, Tx_n) \\
d(Tx_n Tx_{n+1}) &\leq \frac{(a+c+b)}{1-b}d(Tx_{n-1} Tx_n)
\end{aligned} \tag{3.5}$$

Case 2. If $Tx_n \in P, Tx_{n+1} \in Q$ then

$$\begin{aligned}
d(Tx_n, Tx_{n+1}) + d(Tx_{n+1}, y_{n+1}) &= d(Tx_n, y_{n+1}) \\
d(Tx_n, Tx_{n+1}) &\leq d(Tx_n, y_{n+1}) = d(y_n, y_{n+1})
\end{aligned} \tag{3.6}$$

$$\begin{aligned}
d(y_n, y_{n+1}) &= d(Fx_{n-1}, Fx_n) \\
&\leq (a+c)d(Fx_{n-1} Tx_{n-1}) + b[\max\{d(Tx_{n-1}, Fx_{n-1}), d(Tx_{n-1}, Tx_n)\} + d(Tx_n Fx_n)] \\
&= (a+c)d(y_n Tx_{n-1}) + b[\max\{d(Tx_{n-1}, y_n), d(Tx_{n-1} Tx_n)\} + d(Tx_n, y_{n+1})] \\
&= (a+c)d(Tx_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1} Tx_n), d(Tx_{n-1} Tx_n)\}] + b d(Tx_n Tx_{n+1}) \\
&= (a+c)d(Tx_n Tx_{n-1}) + b[\max\{d(Tx_{n-1}, Tx_n), d(Tx_{n-1}, Tx_n)\}] + b d(y_n y_{n+1}) \\
(1-b)d(y_n, y_{n+1}) &\leq (a+c)d(Tx_n, Tx_{n-1}) + b d(Tx_{n-1}, Tx_n) \\
(1-b)d(y_n, y_{n+1}) &\leq (a+c+b)d(Tx_n, Tx_{n-1})
\end{aligned}$$

$$d(y_n, y_{n+1}) \leq \frac{(a+c+b)}{1-b}d(Tx_{n-1}, Tx_n) \tag{3.7}$$

Since $d(Tx_n, Tx_{n+1}) \leq d(y_n y_{n+1})$ from(3.8). Therefore,

$$d(Tx_n, Tx_{n+1}) \leq \frac{(a+c+b)}{1-b}d(Tx_{n-1}, Tx_n) \tag{3.8}$$

Case 3. If $Tx_n \in Q, Tx_{n+1} \in P$ then $Tx_{n-1} \in P$

Since Tx_n is the convex linear combination of Tx_{n-1} and y_n , it follows that

$$d(Tx_n, Tx_{n+1}) \leq \max\{d(Tx_{n-1}, Tx_{n+1}), d(y_n, Tx_{n+1})\}. \tag{3.9}$$

If $d(Tx_{n-1}, Tx_{n+1}) \leq d(y_n, Tx_{n+1})$. Then

$$\begin{aligned}
d(Tx_n, Tx_{n+1}) &\leq d(y_n, Tx_{n+1}) \\
&= d(y_n, y_{n+1}) \\
&= d(Fx_{n-1}, Fx_n) \\
&\leq (a+c)d(Fx_{n-1}, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, Fx_{n-1}), d(Tx_{n-1} Tx_n)\} + d(Tx_n, Fx_n)] \\
&= (a+c)d(y_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, y_n), d(Tx_{n-1}, Tx_n)\} + d(Tx_n, y_{n+1})] \\
&= (a+c)d(y_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, y_n), d(Tx_{n-1}, Tx_n)\}] + b d(Tx_n, Tx_{n+1}) \\
(1-b)d(Tx_n, Tx_{n+1}) &\leq (a+c)d(y_n, Tx_{n-1}) + b[\max\{d(Tx_{n-1}, y_n), d(Tx_{n-1} Tx_n)\}]
\end{aligned} \tag{3.10}$$

Since

$$\begin{aligned}
d(Tx_{n-1}, y_n) + d(y_n, Tx_n) &= d(Tx_{n-1}, Tx_n) \\
d(Tx_{n-1}, y_n) &\leq d(Tx_{n-1}, Tx_n)
\end{aligned}$$

$$\begin{aligned}
(1-b)d(Tx_n, Tx_{n+1}) &\leq (a+c)d(y_n, Tx_{n-1}) + b d(Tx_{n-1}, Tx_n) \\
&\leq (a+c)d(Tx_{n-1}, Tx_n) + b d(Tx_{n-1}, Tx_n) \\
&= (a+c+b) d(Tx_{n-1}, Tx_n) \\
d(Tx_n, Tx_{n+1}) &\leq \frac{(a+c+b)}{1-b} d(Tx_{n-1}, Tx_n). \tag{3.11}
\end{aligned}$$

On the other hand if $d(y_n, Tx_{n+1}) \leq d(Tx_{n-1}, Tx_{n+1})$.

Then $d(Tx_n, Tx_{n+1}) \leq d(Tx_{n-1}, Tx_{n+1}) = d(y_{n-1}, y_{n+1})$ from (3.9).

$$\begin{aligned}
d(Tx_n, Tx_{n+1}) &\leq d(y_{n-1}, y_{n+1}) \\
&= d(Fx_{n-2}, Fx_n) \\
&\leq (a+c)d(Fx_{n-2}, Tx_{n-2}) + b[\max\{d(Tx_{n-2}, Fx_{n-2}), d(Tx_{n-2}, Tx_n)\} + d(Tx_n, Fx_n)] \\
&= (a+c)d(y_{n-1}, Tx_{n-2}) + b[\max\{d(Tx_{n-2}, y_{n-1}), d(Tx_{n-2}, Tx_n)\} + d(Tx_n, y_{n+1})] \\
&= (a+c)d(y_{n-1}, Tx_{n-2}) + b[\max\{d(Tx_{n-2}, y_{n-1}), d(Tx_{n-2}, Tx_n)\} + b d(Tx_n, Tx_{n+1})]
\end{aligned}$$

$$(1-b)d(Tx_n, Tx_{n+1}) \leq (a+c)d(Tx_{n-1}, Tx_{n-2}) + b[\max\{d(Tx_{n-2}, Tx_{n-1}), d(Tx_{n-2}, Tx_n)\}]. \tag{3.12}$$

Since

$$\begin{aligned}
d(Tx_{n-2}, Tx_n) + d(Tx_n, Tx_{n-1}) &= d(Tx_{n-2}, Tx_{n-1}) \\
d(Tx_{n-2}, Tx_n) &\leq d(Tx_{n-2}, Tx_{n-1}).
\end{aligned}$$

Therefore,

$$\begin{aligned}
(1-b)d(Tx_n, Tx_{n+1}) &\leq (a+c)d(Tx_{n-1}, Tx_{n-2}) + b d(Tx_{n-2}, Tx_{n-1}) \\
&= (a+c+b) d(Tx_{n-1}, Tx_{n-2}) \\
d(Tx_n, Tx_{n+1}) &\leq \frac{(a+c+b)}{1-b} d(Tx_{n-1}, Tx_{n-2}). \tag{3.13}
\end{aligned}$$

Thus in the all case we have $d(Tx_n, Tx_{n+1}) \leq A \max\{d(Tx_{n-1}, Tx_n), d(Tx_{n-1}, Tx_{n-2})\}$, where $A = \frac{(a+c+b)}{1-b}$.

It can be easily shown that by induction that for $n \geq 1$ we have

$$d(Tx_n, Tx_{n+1}) \leq A^n \max\{d(Tx_0, Tx_1), d(Tx_1, Tx_2)\}.$$

Now for any positive integer r we have

$$\begin{aligned}
d(Tx_n, Tx_{n+r}) &\leq d(Tx_n, Tx_{n+1}) + d(Tx_{n+1}, Tx_{n+2}) + \dots + d(Tx_{n+r-1}, Tx_{n+r}) \\
&\leq \{1 + A + A^2 + A^3 + \dots + A^{r-1}\} A^n \max\{d(Tx_0, Tx_1), d(Tx_1, Tx_2)\} \\
&\leq \frac{1}{1+A} A^n \max\{d(Tx_0, Tx_1), d(Tx_1, Tx_2)\} \tag{3.14}
\end{aligned}$$

This implies that $d(Tx_n, Tx_{n+1}) \rightarrow 0$ as $n \rightarrow \infty$. So that $\{Tx_n\}$ is a Cauchy sequence and hence converge to a point z in X . We assume that a subsequence $\{Tx_{n_k}\}$ of $\{Tx_n\}$ contained in P and TK is a

closed subspace of X . Since $\{Tx_{n_k}\}$ is Cauchy in TK , it converge to a point, $z \in TK$. Let $u \in T^{-1}z$ then $Tu = z$. Here one also needs to note that $\{Fx_{n_k-1}\}$ will also converge to z .

$$d(Fx_{n_k-1}, Fu) \leq (a+c)d(Fx_{n_k-1}Tx_{n_k-1}) + b[\max\{d(Tx_{n_k-1}, Fx_{n_k-1}), d(Tx_{n_k-1}, Tu)\} + d(Tu, Fu)]$$

Which, on letting, $k \rightarrow \infty$ reduces to

$$d(z, Fu) \leq (a+c)d(z, z) + b \max\{d(z, z), d(z, Tu)\} + bd(Tu, Fu)$$

$$d(Fu, z) \leq bd(Tu, Fu)$$

$$d(Fu, z) \leq bd(Fu, Tu) \tag{3.15}$$

Yielding there by $Tu = Fu$ which shows that u is a point of coincidence for F and T . Since the pair (F, T) is coincidentally commuting, therefore

$$z = Tu = Fu \Rightarrow Fz = FTu = TFu = Tz \tag{3.16}$$

To prove that z is fixed point of F , consider $d(Fz, z) = d(Fz, Fu)$

$$d(Fz, z) \leq (a+b)d(Fz, Tz) + b[\max\{d(Tz, Fz), d(Tz, Tu)\} + d(Tu, Fu)]$$

$$d(Fz, z) \leq (a+b)d(Fz, Tz) + b \max\{d(Tz, Fz), d(Tz, Tu)\} + bd(Tu, Fu)$$

$$d(Fz, z) \leq b d(Fz, z)$$

Which shows that z is a common fixed point of F and T . The proof goes on similar lines in case we assume subsequence $\{Tx_{n_k}\}$ of $\{Tx_n\}$ contained in Q . hence it is omitted. The uniqueness of fixed point follows easily this completes the proof. \square

4 Acknowledgements

The authors would like to express their sincere thanks to the editor and the anonymous referees for their valuable comments and useful suggestions in improving the article.

References

- [1] G.Jungck, *Compatible mappings and common fixed points*, Int. J. Math. Math. Sci., 9(4)(1986), 771–779.
- [2] M.Imdad, *Remarks on fixed point theorems for nonself mappings*, Aligarh Bull. Math., 22(1)(2003), 15–17.
- [3] M.Imdad and S.Kumar, *Boyd and Wong type theorems for two pairs of non-self mappings*, Nonlinear Anal. Forum, 8(1)(2003), 69–78.
- [4] M.D.Khan and R.Bharadwaj, *A fixed point theorem in metrically convex space*, Indian J. Math., 43(3)(2001), 373–379.

- [5] M.S.Khan, H.K.Pathak and M.D.Khan, *Some fixed point theorems in metrically convex spaces*, Georgian Math. J., 7(3)(2000), 523–530.
- [6] M.Imdad, *common fixed point theorems for nonself mapping in metrically convex space via altering Distances*, IJMMS-2005, 24(2005), 4029-4039.
- [7] N.A.Assad, *On a fixed point theorem of Kannan in Banach spaces*, Tamkang J. Math., 7(1)(1976), 91–94.
- [8] N.A.Assad and W.A.Kirk, *Fixed point theorems for set-valued mappings of contractive type*, Pacific J. Math., 43(3)(1972), 553–562.
- [9] O.Had, *On coincidence points in convex metric spaces*, Univ. u Novom Sadu Zb. Rad.Prirod.-Mat. Fak. Ser. Mat., 19(2)(1989), 233–240.
- [10] O.Had and Lj.Gaji, *Coincidence points for set-valued mappings inconvex metric spaces*, Univ. u Novom Sadu Zb. Rad. Prirod.-Mat. Fak. Ser.Mat., 16(1)(1986), 13–25.
- [11] R.Kannan, *Some results on fixed points*, Bull. Calcutta Math. Soc., 60(1968), 71–76.
- [12] R.M.Bianchini, *Su un problema di S. Reich riguardante la teoria dei punti fissi*, Boll. Un. Mat. Ital., 4(5)(1972), 103–108.
- [13] K. Chatterjea, *Fixed-point theorems*, C. R. Acad. Bulgare Sci., 25(1972), 727–730.
- [14] S.Sessa, *On a weak commutativity condition of mappings in fixed point considerations*, Publ. Inst. Math.(Beograd)(N.S.), 32(46)(1982), 149–153.